



Citation for published version:

Mounce, C, Keogh, E & Eccleston, C 2010, 'A principal components analysis of negative affect-related constructs relevant to pain: evidence for a three component structure', *Journal of Pain*, vol. 11, no. 8, pp. 710-717. <https://doi.org/10.1016/j.jpain.2009.10.014>

DOI:

[10.1016/j.jpain.2009.10.014](https://doi.org/10.1016/j.jpain.2009.10.014)

Publication date:

2010

[Link to publication](#)

This is an author's version. A definitive version was subsequently published in *The Journal of Pain*, 11(8), 2010, DOI: 10.1016/j.jpain.2009.10.014

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**A principal components analysis of negative affect-related constructs relevant to
pain: Evidence for a three component structure**

**Charlotte Mounce*, Edmund Keogh, Christopher Eccleston
Centre for Pain Research, University of Bath, UK.**

***Corresponding Author.**

**Address: Centre for Pain Research, The University of Bath, Claverton Down,
Bath, BA2 7AY.**

Tel.: +44 1225 385 434; Fax.: +44 1225 385 434

E-mail address: C.S.Mounce@bath.ac.uk

Keywords: Pain; Factor Analysis; Assessment; Fear; Anxiety

Abstract

A number of negative affect-related constructs are important in pain. Some are general, such as anxiety, depression and negative affectivity, whereas others are more specifically pain-related (e.g., fear of pain, pain anxiety and pain catastrophizing). In addition, some more specific fear-related constructs, such as anxiety sensitivity, illness/injury sensitivity, and fear of negative evaluation, have emerged as important to pain. Although these various constructs are considered conceptually separate, there is likely to be overlap between them. Since the extent of this overlap is unknown, the aim of the current study was to investigate these constructs in one sample in order to identify their common and unique features. Frequently used psychological measures were completed by 508 pain free participants. Principal components analysis resulted in the extraction of three components: 'General distress', 'Fear of pain from injury/insult', and 'Cognitive intrusion of pain'. The results presented here suggests that there is indeed commonality between constructs, which may be due to either an overlap between items within measures or due to close conceptual relatedness. The implications of these core dimensions are discussed with reference to future research and theory.

Perspective

This article explores the relationships between various negative-affect pain-related measures and discusses the results from a principal components analysis. The findings show that some questionnaires may measure the same latent construct. A measure could be developed to measure these three core components more concisely for both clinical and research purposes.

Introduction

Pain and negative affect often co-occur^{5,16} in both clinical and non-clinical presentations of pain. For example, Conrad et al.¹⁰ found that compared with pain free matched controls, chronic pain patients have significantly greater depression and anxiety. This relationship also appears to occur in experimentally induced pain, with depressed mood related to reduced pain tolerance⁴⁸ and higher anxiety to increased pain intensity⁴¹.

Alongside these more general constructs, there are a number of specific anxiety and fear-related dimensions that relate to pain. For example, the pain-related dimensions fear of pain and pain catastrophizing have both been related to pain sensitivity, and both play a role in the development and maintenance of pain-related disability^{3,15,20,27,39,45}. There has also been interest in the fear of anxiety-related sensations (i.e., anxiety sensitivity; AS) and the closely related construct of illness/injury sensitivity (IS)^{14,18,36,43}. Finally, it is becoming increasingly recognised that pain and its associated expression occur within a social context, and so fears related to negative evaluation (FNE; fear of negative evaluation by others) may also be important²⁴.

Although this research indicates that there are both general and specific anxiety-related constructs related to pain, we know less about how they are related to one another. In one of the few studies to examine some of these constructs together, Vancleef et al.⁴⁴ found that IS and AS, but not FNE, were significant predictors of fear of pain and pain-related catastrophizing. However, it remains unclear the degree to which these anxiety-related constructs are distinct from one another, and what the degree of overlap is between them. If there is overlap between these measures, then some constructs may be redundant or less important. Similar observations have been

expressed more generally in the conceptualisation of anxiety and depression, where it has been shown that the scales used to measure these constructs not only need to share a set of similar symptoms, but can be differentiated by examining condition specific characteristics⁴⁶. This possibility has yet to be examined in the context of pain-related anxiety. However, it seems likely that this group of questionnaires measuring constructs relating to negative affect, anxiety, and fear, may have underlying common themes. Indeed, previous research has shown there to be significant correlations between a number of these constructs (e.g. anxiety sensitivity, fear of pain, and pain anxiety⁵⁰).

Our interest here is in the experience of pain and affect in a sample unaffected by a clinical pain state; as such groups are likely to also suffer from co-existing anxiety and depression. Although a non-clinical sample will allow us to examine the naturally occurring relationships between such constructs, our longer term goal is to develop optimal measurement methods of affect in different pain populations. Thus we view the examination of the interrelationship of constructs in a large non-clinical sample as an important and necessary first step, especially given our focus on the function and fidelity of affect constructs and their inter-relatedness. We hypothesized that when these commonly used measures were examined concurrently, in the same population, that significant conceptual overlap would be found, and a higher order affective system would emerge that cuts across independently developed instruments.

Materials and Methods

Participants

The inclusion criteria for participation in this study included self reported good general health and English as a first language. Those reporting chronic pain

(over 3 months) were excluded. An opportunity and snowballing sampling approach was used. For example, students were recruited during lectures, University employees through invitation etc. A total of 508 healthy adults completed the study. Table 1 presents full demographic information for the sample.

TABLE 1

Measures & Procedure

Following Ethics Committee approval from the University of Bath, participants were provided with information about the study. The 508 who provided informed consent to participate were then asked to complete nine questionnaires. The constructs we sought to measure were based upon the constructs found in two models. Firstly, the latest fear avoidance model of chronic pain³⁰ which includes the constructs anxiety sensitivity, negative affectivity, pain catastrophizing, fear of pain, and depression, which are all thought to contribute to the development of chronic pain. Secondly, the hierarchical model of the relationship between negative emotionality, and lower-order constructs²² which has been expanded to include pain relevant constructs¹⁷, which includes negative emotionality (NA), anxiety, illness/injury sensitivity, social evaluation sensitivity (fear of negative evaluation), anxiety sensitivity, pain catastrophizing, and fear of pain. In addition a measure of pain anxiety has been included due to the extensive research in pain on this construct as well as many of the others. The questionnaires were chosen based on theoretical grounds; they are generally viewed as good measures of the underlying constructs under consideration and are judged to be valid and reliable measures¹⁷. Although

some of the measures were developed for use in a chronic pain population, all have been validated and used in a normal population. The constructs sampled (and measures administered) were completed in the following order:

Negative Affect

The Positive and Negative Affect Schedule (PANAS⁴⁷) consists of items relating to ten negative states (e.g. 'distressed', 'upset') and ten positive states (e.g. 'inspired', 'active'). Participants rate the extent to which they feel each affective state on a five point Likert scale (1= very slightly or not at all, 5 = extremely). The participants were asked to answer each item in terms of how they 'generally feel' in order to obtain a trait measure. The 'negative affect' (NA) subscale was used as a measure of negative affectivity. Although a preliminary exploration of the data found that the 'positive affect' subscale formed its own component (as expected), it was not included in the main analyses as the study was more concerned with negative constructs. The NA scale has been found to be highly internally consistent ($\alpha = .84$ to $.87$), have temporal stability over an eight week period ($.71$), and have evidence of convergent ($.81$ to $.92$) and discriminant validity ($.06$ to $-.21$)⁴⁷.

General Trait Anxiety

The Spielberger State/Trait Anxiety Inventory (STAI-t³⁷) was used to measure trait anxiety. The trait form consists of 20 items to which participants indicate the extent these statements apply to them on a four point Likert scale (1 = almost never, 4 = almost always). This scale has been found to have high test-retest coefficients ($.73$ to $.86$ in college students) and internal consistency ($\alpha = .90$), and shows evidence of concurrent validity ($.73$ to $.85$)³⁷.

Depression, Anxiety, and Stress

The short form of the Depression Anxiety and Stress Scale (DASS-21²³) was used as a measure of mood. There are seven items in the subscales relating to depression, anxiety, and stress. The questionnaire is rated on a four point Likert scale (0 = did not apply to me at all, 3 = applied to me very much, or most of the time). Internal consistency ($\alpha = .87$ to $.91$) and concurrent validity ($.55$ to $.85$) was found to be in the acceptable to excellent ranges ². All three subscales were used in the current study.

Fear of illness and injury

The Illness/Injury Sensitivity Index (ISI⁴²) is an 11-item self report which measures fear of illness and injury. Participants are asked how much they agree with each item on a five point Likert scale (1 = very little, 5 = very much). This questionnaire has acceptable internal consistency ($\alpha > .80$)⁴². Factor analysis has revealed two factors, fear of injury and fear of illness⁷. In the present study these two factors were used as subscales.

Catastrophic thinking about pain

The Pain Catastrophizing Scale (PCS³⁸) a 13 item scale, was developed to measure the tendency to catastrophize during painful situations. The items are rated on a five point Likert scale (0 = not at all, 4 = all the time), which serve to form three dimensions: rumination, magnification, and helplessness. The scale has been found to have acceptable test-retest reliability ($.75$), internal consistency ($\alpha = .93$ to $.95$) and predictive validity^{31,32,38}.

Pain Specific Anxiety

The 20-item version of the Pain Anxiety Symptoms Scale (PASS²⁶) was used to measure pain-related anxiety. While it was developed for use with a chronic pain population, it has been used within non-clinical samples, and is considered valid and

reliable in a normal population¹. Items are rated on a 6 point Likert scale (0 = Never, 5 = Always), which are used to form four subscales: fear of pain, escape-avoidance behaviours, physiological symptoms, and cognitive symptoms of anxiety. The PASS-20 has satisfactory internal consistency ($\alpha = .81$ to $.83$) and test-retest reliability (.74 to $.87$), and has good construct and predictive validity^{11,26,35}.

Fear of Pain

The Fear of Pain Questionnaire III (FPQ²⁸) is a 30-item questionnaire in which participants are asked to rate the fear they associate with each item on a five point Likert scale (1 = not at all, 5 = extreme). There are three subscales relating to the fear of minor pain, medical pain, and severe pain. The FPQ has good internal consistency ($\alpha = .87$ to $.88$), test-retest reliability (.69 to $.76$), and predictive validity with high fear of pain individuals as measured by the FPQ having greater avoidance/escape from a behavioural test, compared with low fear of pain individuals²⁸.

Fear of Negative Evaluation

The Brief Fear of Negative Evaluation Scale is a 12 item scale which measures the fear of negative evaluation by others (FNES¹⁹). Items are rated on a five point Likert scale (1 = not at all characteristic of me, 5 = extremely characteristic of me). Individuals who score highly on this questionnaire are more fearful of being negatively evaluated by others. The scale has been found to have good internal consistency ($\alpha = .90$) and test-retest reliability (.75)¹⁹.

Anxiety Sensitivity

The Anxiety Sensitivity Index (ASI³⁴) measures a person's fear associated with the consequences of anxiety-related symptoms. It has 16-items that are rated on a five point Likert scale (1 = very little, 5 = very much). The measure has been found to have good internal consistency ($\alpha = .88$), test-retest reliability (.75) and has been

shown to be a valid measure of the fear of anxiety^{25,33,34}. The ASI is believed to measure three first order factors related to physical concerns, mental incapacitation concerns, and social concerns, which in turn load on to a second order general AS factor⁴⁹. The three subscales were used here.

Statistical approach

The data were analyzed using principal components analyses (PCA) in order to determine whether a smaller number of core components underlie the larger number of psychological constructs related to pain. PCA, like factor analysis, is an exploratory statistical technique used to reduce a large number of variables into a smaller number of subsets. Highly correlated variables which are largely independent of other subsets are combined into components⁴⁰. For more information on this technique see Tabachnick and Fidell⁴⁰. Oblique (oblimin) rotation was used in this case due to the predicted theoretical relationships between these negative constructs. Comrey and Lee⁹ give a guideline for sample size for PCA and suggest 500 to be a “very good” size. Therefore, our sample was perceived to be more than adequate.

Results

Descriptive statistics and correlations

Table 2 presents descriptive statistics for the various measures. Table 3 reports the correlations between total scores, which reveals that they were all positively correlated, and so suitable for principal components analysis. Larger correlations were found within the general mood-related measures (i.e., PANAS-NA, DASS, and STAI-t, FNES), and within the measures related to pain (i.e., PASS, PCS, FPQ, ASI, ISI). This pattern of correlations suggest at least two clusters, one related to mood, the other more specific to pain.

TABLES 2 & 3

Principal Components Analysis

A principal components analysis was carried out on the subscale totals (or the total for those measures without subscales) of the questionnaires using oblique (oblimin) rotation due to potential for correlations between components. In addition, an item analysis was carried out in order to investigate whether the items of subscales load together on the same component. However, primarily the subscale analysis will be reported with reference to the item analysis where necessary. The subscale analysis produced four components with an eigenvalue greater than 1 (factor 1 = 8.2, factor 2 = 2.6, factor 3 = 1.6, factor 4 = 1.1, factor 5 = 0.9), while inspection of the scree plot suggested a three or four component solution. Since the third factor only added a fairly small amount to the variance (12.5%) a two factor solution was also considered. Therefore, two, three, and four components solutions were examined. The two component solution consisted of one component with high loadings from the pain related measures and one with high loadings from general negative mood measures. However, an inspection of the residual correlation matrix showed that a large number of the residuals had values greater than 0.05 suggesting the presence of another factor⁴⁰. This, together with the results of the scree test and eigenvalues resulted in a rejection of a two component solution. Examination of the three and four component solution revealed that the three component solution provided the clearest solution and was therefore selected. The four component solution was very similar with the exception of the two subscales of the ISI forming their own component. In addition it only had an eigenvalue of 1.1. The three component structure is presented in Table 4, and accounted for 59.3% of the total variance. The item PCA closely resembled those

found in the subscale analysis, with the same measures generally loading together in a similar way in both analyses. The only exception to this involved the ISI, which loaded onto 'Fear of pain from injury/insult' on the subscale analysis but in the item analysis they load onto the 'Cognitive intrusion of pain' component. Excluding this, the item analysis and subscale analysis yield very similar solutions. The solution for the subscale PCA accounted for 59.3% of the total variance compared with only 33.9% for the item PCA. It seems likely that this is due to the subscale PCA having a greatly reduced number of variables and more parsimonious solution.

The first component included the subscales from the PASS and the PCS. The PASS and PCS are concerned with cognitions experienced during the experience of pain. Subscales of the PCS are magnification, rumination, and helplessness, all associated with thoughts about pain, specifically catastrophizing about pain. The PASS differs in that it has a subscale devoted to avoidance behaviour and one measuring physiological response to pain, but the other two subscales, fearful thinking and cognitive anxiety, are both concerned with thoughts in response to pain. The highest loading subscales were PASS cognitive anxiety, PCS rumination and PCS helplessness. We also looked at the top loading items from the item-based PCA in order to guide the labelling of the component. This revealed that most of the top-loading items were related to cognitions about pain dominating thoughts. Therefore, this component was named 'cognitive intrusion of pain'. This accounted for 39.2% of the variance.

The second component consisted of the STAI-t total, subscales from the DASS, the FNES total, the PANAS-NA subscale, and the mental concerns subscale of the ASI. The items from these measures are generally related to anxiety, depression, negative affectivity, and fear of negative evaluations from others. Although an

association between depression, negative affectivity, and anxiety has previously been found⁴⁶, the FNES also loaded onto this component suggesting a relationship between these constructs and a concern with what others think. The mental concerns subscale of the ASI also loaded onto this component. The highest loading (sub)scales were STAI-t, DASS depression, and DASS stress. Since the items of the highest loading (sub)scales were related to a mixture of depression, anxiety and stress it seemed that this component was comprised of items measuring general distress and therefore this is the name it was given. This component accounted for 12.5% of the variance.

The third component had the highest loadings from the FPQ subscales, the ASI physical concerns subscales and the ISI subscales. These subscales are principally concerned with the fear of pain that could result from certain circumstances, fear of physical sensations related to anxiety, and fear of illness and injury. The highest loading subscales on this component were the FPQ subscales, the ASI physical concerns subscale and the ISI injury subscale. Examination of the top loading items on the item analysis revealed they related more generally to the fear of various injuries, procedures and physical sensations. This component was named 'fear of pain from injury/insult'. This component accounted for a further 7.6% of the variance. In order to further evaluate the stability of this three component solution, the sample was split into two groups of 254 data sets and a PCA as described above was conducted. Again, the three component solution was selected for each sample due to inspection of eigenvalues, scree plots, and meaningfulness. For the first half of the sample the three components explained 60.8% of the variance and the eigenvalues of these were as follows: factor 1 = 8.8, factor 2 = 2.4, factor 3 = 1.5. For the second half of the sample the three components explained 58.6% of the variance and the eigenvalues were: factor 1 = 7.6, factor 2 = 2.9, factor 3 = 1.8. The three component

solutions for each group closely resemble the solution of the whole sample. The same measures loaded together on the components, with a slight variation of order of component loadings in some places. This finding adds support for the three component solution.

TABLE 4

Component correlations

As previously discussed, the three components were found to be correlated. The largest relationship was found between the two pain related components ‘fear of pain from injury/insult’ and ‘cognitive intrusion of pain’ ($r = .43$). ‘general distress’ was also correlated with these two ($r = .24$, $r = .31$ respectively).

Discussion

The measures used in this study are commonly used in pain research and previous research has examined relationships between some of these constructs. Inspection of correlations found in this study between measures replicates previous findings. For example, measures of anxiety, depression, and negative affectivity had moderate to high correlations ($r = .55$ to $.72$) which is in line with earlier findings⁸. Additionally, moderate correlations were found between trait anxiety and anxiety sensitivity, illness/injury sensitivity, and fear of negative evaluation ($r = .26$ to $.60$) similar to previous research⁴². Finally, correlations found between pain-related constructs fear of pain, anxiety sensitivity, pain catastrophizing, and pain anxiety substantiate previous findings ($r = .46$ to $.71$) (^{44, 50, 4, 38, 12}). Although other studies

have examined relationships between some of these constructs, as far as we are aware, this is the first attempt to consider these measures together. The measures have been developed by different researchers with the aim of measuring a particular aspect of negative mood or fear about pain and conceptually they are thought to reflect different constructs. However, it is likely that there is overlap in what the questionnaires are measuring because they focus on a similar area. Therefore, the aim of this study was to investigate whether anxiety-related constructs, often measured in the context of pain, using instruments with general and overlapping item content, can be reduced to a smaller number of underlying components. When analysed together, three core anxiety constructs emerged: a general affective component ‘general distress’ and two pain-related anxiety/fear components, ‘cognitive intrusion of pain’ and ‘fear of pain from injury/insult’.

With respect to the general distress component, items that loaded on it were related to anxiety, depression, negative affectivity and fear of negative evaluation. It was perhaps unsurprising that these constructs loaded onto the same component as they are conceptually closely related, especially depression, anxiety, and negative affectivity⁴⁶. The other two components were more specifically related to pain. The fear of pain from injury/insult component included items concerned with fear about pain or physical sensations from injury/insult, illness and anxiety, whereas the cognitive intrusion of pain component contained items concerned with cognitive reactions to pain, especially in terms of being unable to take ones thoughts away from the pain.

Closer inspection of the fear of pain from injury/insult and cognitive intrusion of pain components suggests two main differences. The fear of pain from injury/insult component appears to be anticipatory, imagining how one would feel about certain

circumstances, whereas cognitive intrusion of pain is linked to how one reacts to pain. The second difference is that fear of pain from injury/insult is concerned with fear while cognitive intrusion of pain is concerned with anxious thoughts about pain. Although fear and anxiety consist of cognitive, physiological, and behavioural dimensions, fear tends to have a more short-term arousal-like quality associated with it, whereas anxiety has a greater cognitive component, and is less physiologically focused^{3,6}. Therefore, fear of pain from injury/insult may be measuring a more emotional and physiological fear factor, while cognitive intrusion of pain has a more cognitive focus, especially given that the top-loading items tended to relate to intrusive thoughts about pain. Interestingly, other anxiety-related constructs have also made a distinction between emotional and cognitive components (e.g. test anxiety^{21,29}), and so in the same sense the pain questionnaires used here could also be accessing different emotional (fear of pain from injury/insult) and cognitive components (cognitive intrusion of pain).

A second point is that examination of the content of the PASS and PCS reveals that some of it is comparable. This suggests that both measures may be accessing the same (latent) construct that we have labelled ‘cognitive intrusion of pain’. If two questionnaires are being used to measure essentially the same latent construct, then this may have implications in terms of research and for clinical studies, although recognizing that this solution has yet to be replicated with clinical samples. Likewise, inspection of the measures found in fear of pain from injury/insult, the FPQ, ASI, and ISI demonstrates that although the three constructs being measured are related to a different fear: fear of pain from injury, fear of sensations from anxiety symptoms, and a fear of illness/injury, they are also conceptually related, and generally concerned with fear of bodily harm. Therefore, the three questionnaires may

be measuring the same underlying construct ‘fear of pain from injury/insult’.

However, since the questionnaires do appear to be measuring different fears, ‘fear of pain from injury/insult’ may be a higher order factor, with ‘fear of pain’, ‘fear of injury/illness’, and ‘anxiety sensitivity’ being lower order factors in a hierarchical structure. This may be of interest to those researchers who have suggested a hierarchical model of negative emotional²² and pain-related constructs¹⁷.

Interestingly, the three components found here appear in one form or another within the fear avoidance model⁴⁵ which suggests that negative affectivity, pain catastrophizing, and pain-related fear are integral in contributing to the development and maintenance of chronic pain. Amended versions have included anxiety sensitivity³⁰ and pain-related anxiety, although it appears to be portrayed as separate to pain catastrophizing³. The results of the current study suggest that there may be overlap between questionnaires that are used to measure pain-related anxiety and pain catastrophizing (the PASS and PCS), and so the fear avoidance model may need to take this into consideration. Indeed, although fear of pain has been suggested to be critical in the development of chronic pain and pain-related disability²⁰, less is known about whether illness/injury sensitivity and anxiety sensitivity contribute to the development and maintenance of pain and disability. The results of this study suggest that the three constructs are closely related and so may also serve as vulnerability factors for the development of chronic pain. Therefore, a fear of anxiety symptoms and illness/injury may need to also be considered with respect to the fear-avoidance model and may also be an important area of focus in a clinical setting.

In another study to consider a number of self-report measures relevant to pain, Davidson et al (2008)¹³ carried out an exploratory factor analysis (EFA) on a number of questionnaires including those measuring sensory pain, functioning, coping with

pain, and pain disability in patients with chronic pain. Seven factors were extracted. Measures of anxiety and depression loaded together, but the Pain Catastrophizing Scale also loaded onto this factor. However, since this study was examining a wide range of constructs measuring the different aspects of pain experience, it seems likely that when loaded into an EFA the Pain Catastrophizing Scale may load with depression and anxiety as other measures of negative emotions and cognitions. Since the present study only examined negative emotions and cognitions related to pain, this allowed a more detailed understanding of how these constructs related to each other.

Implications of this study are that some questionnaires used in pain measurement often have overlapping content and therefore one or more may be redundant. In addition, the findings contribute to the theory of the hierarchical model of negative and pain-related constructs^{22,17}. Further, the relationships found between the constructs may be of interest in terms of adding additional constructs or deleting certain constructs from the fear-avoidance model^{3,20}. Finally, the findings of this study can be used in the development of a brief questionnaire to reduce the number of measures necessary.

The study has some relevant limitations. First, we were interested in how the various constructs under consideration were related within a healthy sample i.e., in absence of chronic pain. It is entirely possible that an alternative structure may be found within chronic pain patients and extrapolation to clinical samples should be done with caution. Related to this is the idea that more specific constructs (e.g., worry about pain) may only become relevant in a chronic pain context as it is largely defined by the existence of repetitive threat. Additionally, although we excluded those with chronic pain, the experiences of pain of the sample in this study are likely to vary widely and include some participants who are familiar with pain. However, since

chronic pain was an exclusion criterion and 70 percent of the sample was under the age of 35, this is likely to be restricted. However, this needs to be taken into consideration when interpreting the results. Second, although we included nine instruments, others were not selected and could offer different solutions. We cannot, therefore, definitively state that we have sampled all relevant content. Third, PCA itself can be criticized as it has no available criteria against which to test the solution, and since following extraction there is an infinite number of rotations available. The final decision and labelling depends upon expert decision and interpretation⁴⁰.

Future research could focus on further analysis of the stability of this factor structure with other healthy samples, and samples of patients with chronic pain. If this tripartite solution is robustly replicated a stronger argument can be made for the development of a brief questionnaire to directly measure these three core components economically, increasing accuracy and reducing demand on participants.

Acknowledgement

This work was funded by a University Research Studentship from the Psychology Department, University of Bath.

References

1. Abrams MP, Carleton RN, Asmundson GJG: An exploration of the psychometric properties of the PASS-20 with a nonclinical sample. *J Pain*: 8(11), 879-886, 2007
2. Antony MM, Bieling PJ, Cox BJ, Enns MW, Swinson RP: Psychometric Properties of the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample. *Psychol Assessment*: 10(2), 176-181, 1998
3. Asmundson GJ, Norton PJ, Vlaeyen JWS: Fear-avoidance models of chronic pain: An overview. In: *Understanding and treating fear of pain*, (Asmundson, G.J., Vlaeyen, J. W. S., and Crombez, G., Eds.), Oxford University Press, Oxford, 2004, pp. 3-24.
4. Asmundson GJG, and Taylor S: Role of anxiety sensitivity in pain-related fear and avoidance. *J Behav Med*, 19(6), 577-586, 1996
5. Banks SM, Kerns RD: Explaining high rates of depression in chronic pain: a diathesis-stress framework. *Psychol Bull*: 119(1), 95-110, 1996
6. *Anxiety and its Disorders*, 2nd edition, Barlow DH, Guilford Press, New York, 2002.
7. Carleton RN, Asmundson GJ, Taylor S: Fear of physical harm: Factor structure and psychometric properties of the injury/illness sensitivity index. *J Psychopathol Behav*: 27(4), 235-241, 2005
8. Clark LA, and Watson D: Tripartite model of anxiety and depression: Psychometric evidence and taxonomic implications. *J Abnorm Psychol*, 100, 316-336, 1991
9. Comrey, A. L., & Lee, H. B. (1992). *A First Course in Factor Analysis* (2nd ed.). Hillsdale, NJ: Lawrence Erlbaum Associates.

10. Conrad R, Schilling G, Bausch C, Nadstawek J, Wartenberg HC, Wegener I, Geiser F, Imbierowicz K, Liedtke R: Temperament and character personality profiles and personality disorders in chronic pain patients. *Pain*: 133, 197-209, 2007
11. Coons MJ, Hadjistavropoulos HD, Asmundson GJ: Factor structure and psychometric properties of the Pain Anxiety Symptoms Scale-20 in a community physiotherapy clinic sample. *Eur J Pain*: 8, 511-516, 2004
12. Crombez G, Vlaeyen JWS, Heuts PHTG, and Lysens R: Pain-related fear is more disabling than fear itself: evidence on the role of pain-related fear in chronic back pain disability. *Pain*, 80, 329-339, 1999.
13. Davidson MA, Tripp DA, Fabrigar LR, Davidson PR: Chronic pain assessment: A seven-factor model. *Pain Res Manage*: 13(4), 299-308, 2008.
14. Drahovzal DN, Stewart SH, Sullivan MJL: Tendency to catastrophize somatic sensations: Pain catastrophizing and anxiety sensitivity in predicting headache. *Cogn Behav Ther*: 35(4), 226-235, 2006
15. Keefe FJ, Brown GK, Wallston KA: Coping with rheumatoid arthritis: catastrophizing as a maladaptive strategy. *Pain*: 37, 51-56, 1989
16. Keefe FJ, Lumley M, Anderson T, Lynch T, Carson KL: Pain and emotion: New research directions. *J Clin Psychol*: 57, 587-607, 2001
17. Keogh E, Asmundson GJ: Negative affectivity, catastrophizing, and anxiety sensitivity. In: *Understanding and Treating Fear of Pain* (Asmundson, G. J., Vlaeyen, J. W. S. and Crombez, G. Eds.), Oxford University Press, Oxford, 2004, pp. 91-115.

18. Keogh E, Barlow C, Mounce C, Bond FW: Assessing the relationship between cold pressor pain responses and dimensions of the anxiety sensitivity profile in healthy men and women. *Cogn Behav Ther*: 35(4), 198-206, 2006
19. Leary MR: A brief version of the fear of negative evaluation scale. *Pers Soc Psychol B*: 9(3), 371-375, 1983
20. Leeuw M, Goossens MEJB, Linton SJ, Crombez G, Boersma K, Vlaeyen JWS: The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med*: 30(1), 77-94, 2007
21. Liebert RM, Morris LW: Cognitive and emotional components of test anxiety: a distinction and some initial data. *Psychol Rep*: 20, 975-978, 1967
22. Lilienfeld SO, Turner SM, Jacob RG: Anxiety sensitivity: An examination of theoretical and methodological issues. *Adv Behav Res Ther*: 15, 147-183, 1993
23. Manual for Depression Anxiety Stress Scales, 2nd edition, Lovibond SH, Lovibond PF, Psychology Foundation, Sydney, 1995.
24. MacDonald G, Leary MR: Why does social exclusion hurt? The relationship between social and physical pain. *Psychol Bull*: 131, 202-223, 2005
25. Maller RG, Reiss S: A behavioral validation of the Anxiety Sensitivity Index. *J Anxiety Disord*: 1, 265-272, 1987
26. McCracken LM, Dhingra L: A short version of the pain anxiety symptoms scale (PASS-20): Preliminary development and validity. *Pain Res Manag*: 7(1), 45-50, 2002
27. McCracken LM, Zayfert C, Gross RT: The Pain Anxiety Symptoms Scale: Development and validation of a scale to measure fear of pain. *Pain*: 50, 67-73, 1992

28. McNeil DW, Rainwater AJ: Development of the Fear of Pain questionnaire - III. J Behav Med: 21(4), 389-410, 1998
29. Morris LW, Davis MA, Hutchings CH: Cognitive and emotional components of anxiety: literature review and a revised worry-emotionality scale. J Educ Psychol: 73, 541-555, 1981
30. Norton PJ, Asmundson GJ: Amending the fear-avoidance model of chronic pain: What is the role of psychological arousal? Behav Ther: 34, 17-30, 2003
31. Osman A, Barrios FX, Gutierrez PM, Kopper BA, Merrifield T, Grittmann L: The Pain Catastrophizing Scale: Further psychometric evaluation with adult samples. J Behav Med: 23(4), 351-365, 2000
32. Osman A, Barrios FX, Kopper BA, Hauptmann W, Jones J, O'Neill E: Factor structure, reliability, and validity of the Pain Catastrophizing Scale. J Behav Med: 20(6), 589-605, 1997
33. Peterson RA, Heilbronner RL: The Anxiety Sensitivity Index: Construct validity and factor analytic structure. J Anx Disord: 1(2), 117-121, 1987
34. Reiss S, Peterson RA, Gursky DM, McNally RJ: Anxiety Sensitivity, Anxiety Frequency and the Prediction of Fearfulness. Behav Res Ther: 24(1), 1-8, 1986
35. Roelofs J, McCracken LM, Peters M L, Crombez G, van Breukelen G, Vlaeyen JWS: Psychometric evaluation of the Pain Anxiety Symptoms Scale (PASS) in chronic pain patients. J Behav Med: 27(2), 167-183, 2004
36. Schmidt NB, Cook JH: Effects of anxiety sensitivity on anxiety and pain during a cold pressor challenge in patients with panic disorder. Behav Res Ther: 37, 313-323, 1999

37. Manual for the State-trait Anxiety Inventory (Form Y), Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA, Consulting Psychology Press, Palo Alto, CA, 1983.
38. Sullivan MJL, Bishop SR, Pivik J: The Pain Catastrophizing Scale: Development and validation. *Psychol Assess*: 7(4), 524-532, 1995
39. Sullivan MJL, Thorn B, Haythornthwaite JA, Keefe FJ, Martin M, Bradley LA, Lefebvre JC: Theoretical perspectives on the relationship between catastrophizing and pain. *Clin J Pain*: 17, 52-64, 2001
40. Tabachnick BG, Fidell LS: *Using Multivariate Statistics*, 5th edition, Pearson Education Inc, Boston, 2007.
41. Tang J, Gibson SJ: A psychophysical evaluation of the relationship between trait anxiety, pain perception, and induced state anxiety. *J Pain*: 6(9), 612-619, 2005
42. Taylor S: The structure of fundamental fears. *J Behav Ther Exp Psy*: 24(4), 289-299, 1993
43. Vancleef LMG, Peters ML, Gilissen SMP, De Jong PJ: Understanding the role of injury/illness sensitivity and anxiety sensitivity in (automatic) pain processing: An examination using the extrinsic affective simon task. *J Pain*: 8, 563-572, 2007
44. Vancleef LMG, Peters ML, Roelofs J, Asmundson GJ: Do fundamental fears differentially contribute to pain-related fear and pain catastrophizing? An evaluation of the sensitivity index. *Eur J Pain*: 10, 527-536, 2006
45. Vlaeyen JWS, Linton SJ: Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain*: 85, 317-322, 2000

46. Watson D, Clark AC: Affects seperable and inseperable: On the hierarchical arrangements of the negative effects. *J Pers Soc Psychol*: 62, 489-505, 1992
47. Watson D, Clark AC, Tellegen A: Development and validation of brief measures of positive and negative effect: The PANAS Scales. *J Pers Soc Psychol*: 54, 1063-1070, 1988
48. Zelman DC, Howland EW, Nichols SN, Cleeland CS: The effects of induced mood on laboratory pain. *Pain*: 46, 105-111, 1991
49. Zinbarg RE, Barlow DH, Brown TA: Hierarchical structure and general factor saturation of the Anxiety Sensitivity Index: Evidence and implications. *Psychol Assess*: 9, 277-284, 1997
50. Zvolensky MJ, Goodie JL, McNeil DW, Sperry JA, Sorrell JT: Anxiety sensitivity in the prediction of pain-related fear and anxiety in a heterogeneous chronic pain population. *Behav Res Ther*: 39, 683-696, 2001

Table 1. Demographic details of the sample.

	<i>Mean or Proportion</i>
Age	
Male	34.9 (SD 12.3)
Female	34.7 (SD 14.3)
Sex	
Male	38.6%
Female	59.3%
Missing	2.2%
Occupation	
Students	17.7%
Non-manual	65.5%
Manual (skilled and unskilled)	12.0%
Missing	4.7%
Ethnic group	
White European	92.7%
Other ethnicities	4.6%
Missing	2.8%

Table 2. Descriptive statistics for the various questionnaire totals and subscales.

	Mean total score	Standard deviation	Range
FNES	34.87	10.05	12-60
PANAS-NA	17.06	5.67	10-50
STAI-t	38.66	9.13	20-80
DASS total	10.15	8.15	0-63
DASS depression	2.83	3.16	0-21
DASS anxiety	1.99	2.54	0-21
DASS stress	5.34	3.97	0-21
ISI total	23.61	8.77	11-55
ISI illness	16.01	5.74	6-30
ISI injury	7.60	3.59	5-25
PCS total	12.00	9.09	0-52
PCS rumination	5.07	3.82	0-16
PCS magnification	2.29	2.07	0-12
PCS helplessness	4.64	4.17	0-24
PASS total	22.22	15.65	0-100
PASS fearful appraisal	3.55	3.86	0-25
PASS cognitive anxiety	7.55	5.14	0-25
PASS physiological anxiety	3.87	3.96	0-25
PASS escape and avoidance behaviour	7.25	5.19	0-25
FPQ total	66.96	19.95	30-150

FPQ minor	15.73	5.83	10-50
FPQ severe	29.73	9.66	10-50
FPQ medical	21.50	7.85	10-50
ASI total	29.81	8.41	16-80
ASI physical concerns	14.42	5.26	8-40
ASI mental concerns	5.61	2.17	4-20
ASI social concerns	9.79	2.74	4-20

FNES = Fear of Negative Evaluation Scale, PANAS-NA = Positive and Negative Affectivity Scale – negative affect subscale, STAI-t = Spielberger State-Trait Anxiety Inventory – Trait version, DASS = Depression, Anxiety, Stress Scale, ISI = Injury/Illness Sensitivity Index, PCS = Pain Catastrophizing Scale, PASS = Pain Anxiety Symptoms Scale, FPQ = Fear of Pain Questionnaire, ASI = Anxiety Sensitivity Index.

Table 3: Correlations between total scores of each questionnaire

	NA	STAI-t	DASS	ISI	PCS	PASS	FPQ	ASI
FNES	.39	.60	.49	.24	.26	.28	.30	.40
NA		.64	.55	.29	.32	.31	.24	.36
STAI-t			.72	.26	.29	.31	.26	.41
DASS				.23	.30	.33	.21	.42
ISI					.48	.46	.49	.48
PCS						.71	.46	.51
PASS							.49	.54
FPQ								.59

Note. All correlations are significant at the $p < 0.01$ level. FNES = Fear of Negative Evaluation Scale, NA = negative affect subscale of the Positive and Negative Affectivity Scale, STAI-t = Spielberger State-Trait Anxiety Inventory – trait version, DASS = Depression, Anxiety, Stress Scale, ISI = Injury/Illness Sensitivity Index, PCS = Pain Catastrophizing Scale, PASS = Pain Anxiety Symptoms Scale, FPQ = Fear of Pain Questionnaire, ASI = Anxiety Sensitivity Index.

Table 4. Factor loadings of subscale totals.

Subscales	Cognitive intrusion of pain	General distress	Fear of pain from injury/insult
PASS cognitive anxiety	.91	.01	-.04
PCS rumination	.82	-.04	.02
PCS helplessness	.79	.01	.05
PASS escape and avoidance behaviors	.79	-.06	-.06
PASS fearful appraisal of pain	.76	.08	.09
PCS magnification	.75	.08	.06
PASS physiological anxiety	.71	.11	.06
STAI-t	-.01	.87	.02
DASS depression	.06	.82	-.19
DASS stress	.04	.81	-.08
DASS anxiety	.03	.75	.02
PANAS – NA	.09	.67	.01
FNES	-.07	.62	.19
ASI mental concerns	.05	.46	.31
FPQ severe pain	.05	-.10	.80
FPQ minor pain	-.03	-.02	.79
FPQ medical pain	-.03	.01	.78
ASI physical concerns	.20	.13	.60
ISI injury	.16	-.10	.59

ISI illness	.23	.04	.56
ASI social concerns	-.07	.27	.52

Note. FNES = Fear of Negative Evaluation Scale, NA = negative affect subscale of the Positive and Negative Affectivity Scale, STAI-t = Spielberger State-Trait Anxiety Inventory - trait version, DASS = Depression, Anxiety, Stress Scale, ISI = Injury/Illness Sensitivity Index, PCS = Pain Catastrophizing Scale, PASS = Pain Anxiety Symptoms Scale, FPQ = Fear of Pain Questionnaire, ASI = Anxiety Sensitivity Index. Salient loadings ($\geq .3$) appear in **bold**.